

Rehydrat®

agilan® complementary feed

To support digestion

Highly soluble, effervescent, rich in energy and electrolytes

For calves, piglets and poultry

Composition

Lactose, Sodium bicarbonate, Sodium chloride, Potassium chloride

Analytical constituents

Crude protein	5.4 %
Crude fat	< 0.3 %
Crude fiber	< 0.5 %
Crude ash	13.0 %
Sodium	6.0 %
Potassium	1.3 %
Chloride	4.1 %
Lysine	0.0 %
Methionine	0.0 %

Content of additives per kg

Vitamins: Vitamin C (3a300) 10,000 mg

Recommended use

For oral rehydration in *calves, piglets and poultry*.

Rehydrat® quickly and reliably compensates electrolyte and body water losses due to an adequate mineral composition and reverses energy deficiencies due to a high content of energy-rich lactose.

Feeding recommendation

Calves: 100 g/2 liters of lukewarm water

should be administered in addition to the usual milk feeds as 1 - 3 feeds per day in a daily amount corresponding to the severity of the cases for independent intake.

The Rehydrat® solution should be used for the entire duration of an altered fecal consistency.

Fecal consistency	Additional liquid requirement (in % of b.w.)	40 kg calf	
		per day	waterings per day
Thin mushy	approx. 5 %	2 l	2 l / 1 x daily
Soupy	approx. 10 %	4 l	2 l / 2 x daily
Watery	15 (-20) %	6 (-8 l)	2 l / 3 x daily

Piglets: 100 g/2 liters of lukewarm water

The Rehydrat® solution is offered to the piglets for autonomous intake. It is recommended to freshly prepare the solution several times a day. This maintains the piglets' curiosity and prevents odor absorption from the environment.

Poultry: Dissolve 100 g/200 liters of drinking water.



Rehydrat®

Feeding instruction

The solution should be freshly prepared with clean water prior to each use. Rehydrat® should be administered at intervals to the ration of whole milk / milk replacer. Due to lactose Rehydrat® is very rich in energy with a low osmolarity. Lactose maintains the lactase activity so that it is possible to return to milk feeding without any problems. This product must not be used as a sole drinker for calves and piglets.

Recommended feeding period: 1 - 7 days.

In case of risk of, during periods of or after digestive disturbance. It is recommended that a veterinarian's opinion be sought before use or before extending the feeding period.

Storage

Shelf life guarantee is only valid for unopened packages, stored at a cool and dry place. Keep the product out of the reach of children!

Date of revision of the text: December 2022