

Tricho-Stop® Powder

agilan® Complementary feed

For all animal species

Composition

Wheat semolina bran, sodium chloride, malt rootlets, brewer's yeast, calcium carbonate

Analytical constituents

Crude protein	13.5 %	Crude fibre	6.5 %
Crude fat	3.0 %	Crude ash	33.0 %
Sodium	9.0 %	Lysine	0.7 %
Methionine	0.2 %		

Recommended use

For support of the physiological skin flora in nutrition-related disorders.

Content of additives per kg

Trace elements:

Zinc (3b605) 11,000 mg

(in form of zinc sulphate monohydrate)

Iron (3b103) 2,000 mg

(in form of iron(II)sulphate monohydrate)

Copper (3b405) 350 mg

(in form of copper(II)sulphate pentahydrate)

Feeding recommendation

Calves: up to 35 g / animal and day (½ measuring spoon)

Cows, bulls, heifers and horses: up to 70 g / animal and day (1 measuring spoon)

Other animals (e.g. poultry): 7 – 10 kg/ton of feed

Feed at least for a period of 30 days.

Feeding instructions

Due to the increased content of trace elements compared to complete feed, this complementary feed may only be used up to max. 0,8 % of the daily ration. The maximum content (total) of zinc, copper and iron in the complete feed must be taken into account. Feedstuff like malt rootlets may contain the following substance(s): Hordenine

Storage

Shelf life guarantee is only valid for unopened packages, stored at a cool and dry place.

Date of revision of the text: May 2022

6853/K/05/22-eng

